

# Rewilding - det manglende led i naturgenopretningen?



Jens-Christian Svenning, professor, centerleder, VILLUM Investigator  
*Department of Bioscience - Aarhus University*

# General framework for rewilding as a restoration approach

- Rewilding
  - **Ecological restoration to promote self-regulating biodiverse (complex) ecosystems**
  - Key aspects
    - Reducing human control
    - Restoring ecological integrity (natural processes)
  - **Design and implementation**

RESEARCH

## REVIEW SUMMARY

### REWILDING

## Rewilding complex ecosystems

Andrea Perino\*, Henrique M. Pereira\*, Laetitia M. Navarro, Néstor Fernández, James M. Bullock, Silvia Ceaușu, Ainara Cortés-Avizanda, Roel van Klink, Tobias Kuehmerle, Angela Lomba, Guy Pe'er, Tobias Plieninger, José M. Rey Benayas, Christopher J. Sandom, Jens-Christian Svenning, Helen C. Wheeler

**BACKGROUND:** Rapid global change is creating fundamental challenges for the persistence of natural ecosystems and their biodiversity. Conservation efforts aimed at the protection of landscapes have had mixed success, and there is an increasing awareness that the long-term protection of biodiversity requires inclusion of flexible restoration along with protection. Rewilding is one such approach that has been both promoted and criticized in recent years. Proponents emphasize the potential of rewilding to tap opportunities for restoration while creating benefits for both ecosystems and societies. Critics discuss the lack of a consistent definition of rewilding and insufficient knowledge about its potential outcomes. Other criticisms arise from the mistaken notion that rewilding actions are planned without considering societal acceptability and benefits. Here, we present a framework for rewilding actions that can serve as a guideline for researchers and managers. The framework is applicable to a variety of rewilding approaches, ranging from passive to trophic rewilding, and aims

to promote beneficial interactions between society and nature.


**ADVANCES:** The concept of rewilding has evolved from its initial emphasis on protecting large, connected areas for large carnivore conservation to a process-oriented, dynamic approach. On the basis of concepts from resilience and complexity theory of social-ecological systems, we identify trophic complexity, stochastic disturbances, and dispersal as three critical components of natural ecosystem dynamics. We propose that the restoration of these processes, and their interactions, can lead to increased self-sustainability of ecosystems and should be at the core of rewilding actions. Building on these concepts, we develop a framework to design and evaluate rewilding plans. Alongside ecological restoration goals, our framework emphasizes people's perceptions and experiences of wildness and the regulating and material contributions from restoring nature. These societal aspects are important outcomes and may be critical factors for the success of

rewilding initiatives (see the figure). We further identify current societal constraints on rewilding and suggest actions to mitigate them.

**OUTLOOK:** The concept of rewilding challenges us to rethink the way we manage nature and to broaden our vision about how nature will respond to changes that society brings, both intentionally and unintentionally. The effects of rewilding actions will be specific to each ecosystem, and thus a deep understanding of the processes that shape ecosystems is critical to anticipate these effects and to take appropriate management actions. In addition, the decision of whether a rewilding approach is desirable should consider stakeholders' needs and expectations. To this end, structured restoration planning—based on participatory processes involving researchers, managers, and stakeholders—that includes monitoring and adaptive management can be used. With the recent designation of 2021–2030 as the “decade of ecosystem restoration” by the United Nations General Assembly, policy- and decision-makers could push rewilding topics to the forefront of discussions about how to reach post-2020 biodiversity goals. ■

The list of author affiliations is available in the full article online.  
\*Corresponding author. Email: andrea.perino@idee.de (A.P.); hpereira@idee.de (H.M.P.)  
Cite this article as: Perino et al., *Science* 364, eaav5570 (2019). DOI: 10.1126/science.aav5570

**TOMORROW'S EARTH**  
Read more articles online at [scim.ag/tomorrows-earth](http://scim.ag/tomorrows-earth)



**Rewilding actions and outcomes are framed by societal and ecological context.** Rewilding can be assessed by representing the state of ecosystems in a three-dimensional space where each dimension corresponds to an ecological process. The difference in volume between the restored (yellow pyramid) and the degraded ecosystem (orange pyramid) is a proxy for the effects of rewilding on the self-sustainability of

the ecosystem. The dashed line within the yellow pyramid represents the societal boundaries that determine to what extent ecological processes can be restored. Rewilding actions can help push societal boundaries toward the ecological potential (orange arrows) by promoting societal support and opportunities for people to experience the autonomy of ecological processes in enjoyable ways.

Perino et al., *Science* 364, 351 (2019) | 26 April 2019


1 of 1

# Three key ecological components

- Trophic complexity
- Natural disturbances
- Connectivity/Dispersal

## Ecological state

Restoration of ecological processes can positively influence their interactions – e.g., species diversity and trophic complexity can be increased if dispersal to new ecosystems is possible.



# Trophic rewilding

- Definition
  - **Species introductions to restore top-down trophic interactions and associated trophic cascades to promote self-regulating biodiverse ecosystems** (Svenning et al. 2016 PNAS)
- Mostly megafauna-based, due to
  - Ecological importance
  - Size-biased defaunation

SPECIAL FEATURE  
PERSPECTIVE



SPECIAL FEATURE: PERSPECTIVE

## Science for a wilder Anthropocene: Synthesis and future directions for trophic rewilding research

Jens-Christian Svenning<sup>a,1,2</sup>, Pål B. M. Pedersen<sup>a,1</sup>, C. Josh Donlan<sup>b,c</sup>, Rasmus Ejrnæs<sup>d</sup>, Søren Faurby<sup>e</sup>, Mauro Galetti<sup>f</sup>, Dennis M. Hansen<sup>g</sup>, Brody Sandel<sup>h</sup>, Christopher J. Sandom<sup>h</sup>, John W. Terborgh<sup>i</sup>, and Frans W. M. Vera<sup>j</sup>

Edited by Yadvinder Malhi, Oxford University, Oxford, United Kingdom, and accepted by the Editorial Board August 5, 2015 (received for review March 16, 2015)

Trophic rewilding is an ecological restoration strategy that uses species introductions to restore top-down trophic interactions and associated trophic cascades to promote self-regulating biodiverse ecosystems. Given the importance of large animals in trophic cascades and their widespread losses and resulting trophic downgrading, it often focuses on restoring functional megafaunas. Trophic rewilding is increasingly being implemented for conservation, but remains controversial. Here, we provide a synthesis of its current scientific basis, highlighting trophic cascades as the key conceptual framework, discussing the main lessons learned from ongoing rewilding projects, systematically reviewing the current literature, and highlighting unintentional rewilding and spontaneous wildlife comebacks as underused sources of information. Together, these lines of evidence show that trophic cascades may be restored via species reintroductions and ecological replacements. It is clear, however, that megafauna effects may be affected by poorly understood trophic complexity effects and interactions with landscape settings, human activities, and other factors. Unfortunately, empirical research on trophic rewilding is still rare, fragmented, and geographically biased, with the literature dominated by essays and opinion pieces. We highlight the need for applied programs to include hypothesis testing and science-based monitoring, and outline priorities for future research, notably assessing the role of trophic complexity, interplay with landscape settings, land use, and climate change, as well as developing the global scope for rewilding and tools to optimize benefits and reduce human-wildlife conflicts. Finally, we recommend developing a decision framework for species selection, building on functional and phylogenetic information and with attention to the potential contribution from synthetic biology.

conservation | megafauna | reintroduction | restoration | trophic cascades

Human impacts are so pervasive that a new geological epoch has been proposed: the Anthropocene (1). The effects on ecosystems and biodiversity are one of the biggest challenges facing modern society. Large-bodied animals are particularly affected, with massive prehistoric extinctions (2–4) and severe declines in many extant species (5). Over the last decades it has become increasingly clear that large animals are often important for ecosystem function and biodiversity via

trophic cascades, the propagation of consumer impacts downward through food webs (6, 7). Their widespread losses have led to trophic downgrading on a global scale, with negative effects on ecosystems and biodiversity (6–8).

These observations have inspired a new ecological restoration approach that we here refer to as “trophic rewilding.” The rewilding concept was introduced in the late 20th century as a large-scale conservation

<sup>a</sup>Section for Ecoinformatics & Biodiversity, Department of Bioscience, Aarhus University, DK-8000 Aarhus C, Denmark; <sup>b</sup>Advanced Conservation Strategies, Midway, UT 84049; <sup>c</sup>Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, NY 14853; <sup>d</sup>Section for Biodiversity & Conservation, Department of Bioscience, Aarhus University, DK-8410 Randø, Denmark; <sup>e</sup>Departamento de Ecología, Universidade Estadual Paulista, 13506-900 Rio Claro, São Paulo, Brazil; <sup>f</sup>Institute of Evolutionary Biology and Environmental Studies, University of Zurich, 8057 Zurich, Switzerland; <sup>g</sup>Wildlife Conservation Research Unit, Department of Zoology, Oxford University, The Recanati-Kaplan Centre, Oxfordshire OX13 5QL, United Kingdom; <sup>h</sup>Center for Tropical Conservation, Nicholas School of the Environment and Earth Sciences, Duke University, Durham, NC 27708; and <sup>i</sup>Community and Conservation Ecology, Groningen Institute for Evolutionary Life Sciences, University of Groningen, Cocoon, 9700 CC Groningen, The Netherlands

Author contributions: J.-C.S., P.B.M.P., and S.F. designed research; J.-C.S., P.B.M.P., S.F., and D.M.H. performed research; P.B.M.P. and S.F. analyzed data; and J.-C.S., P.B.M.P., C.J.D., R.E., S.F., M.G., D.M.H., B.S., C.J.S., J.W.T., and F.W.M.V. wrote the paper. The authors declare no conflict of interest.


This article is a PNAS Direct Submission. Y.M. is a guest editor invited by the Editorial Board.

<sup>1</sup>J.-C.S. and P.B.M.P. contributed equally to this work.

<sup>2</sup>To whom correspondence should be addressed. Email: svenning@bios.au.dk.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1502556112/-DCS/supplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1502556112/-DCS/supplemental).

# Idea: Megafauna promotes biodiversity via top-down trophic processes+



**Increase diversity capacity of natural and semi-natural areas**

# Abiotic & passive rewilding

- Abiotic rewilding
  - Restoration of natural physical processes
- Passive rewilding  
(=Passive management)
  - Spontaneous ecological dynamics without any, even initial management
  - Always an important aspect



Marselisskovene



Vorsø, Horsens Fjord

# Why is rewilding needed?


- 1) Evolutionary perspective on biodiversity
- 2) Wildness as a value
- 3) Overcome shifting baselines
- 4) Scaling
- 5) Dynamic

# **1: EVOLUTIONARY PERSPECTIVE**


# Current species are ancient = evolved & persistent in wild ecosystems



Cultural landscape



Million years ago



Extant species

Mammals Beetles Trees Vascular plants



*Evolutionary background of ecological adaptations even deeper*


# Skylark (*Alauda arvensis*)



# Beech (*Fagus sylvatica*)



# Current species diversity evolved in megafauna-rich ecosystems




**Rich megafaunas the evolutionary norm  
(an evolutionary base-line)**

Rich megafaunas have been the  
standard for millions of years



England (Trafalgar Square) 125,000

Rich megafaunas have been the standard for millions of years



Germany (Dorn-Dürkheim), 8 million years ago (Wolfgang Weber)



De store dyr er vigtige

Frahegning i Yellowstone National Park

# De store dyr er vigtige


**Tabel 4.2.** Oversigtstabel over faglige skøn og vurderinger af biodiversitetseffekter ved forskellige plejemetoder. Jo flere +’er, jo større positiv effekt.

Type af effekt	Maskinel biomassehøst	Intensiv sommer- græsning	Sommer- græsning	Rotations- græsning	Vinter- græsning	Helårs- græsning	Vild- græsning
Hæmme tilgroning	+++	++	+	+	++	++	++
Hæmme konkurrenceplanter	+++	+++	++	++	+	++	+
Skabe blottet jord	++	+	-/+	+	+++	+++	+++
Abiotisk variation	-/++	++	++	++	+++	+++	++++
Sprede frø	-	+	+	+	+	++	++++
Kulstof variation	-/++	+	+	++	++	+++	++++
Blomstring	++	-	-/+	++	++	+++	+++
Hvirvelløse dyr	-	-	-/+	+	++	+++	+++
Lysstillede veterantræer	+	++	+++	+++	+++	+++	+++
Gødning til faunaen	-	++	++	+	-	+++	++++
Ådsler	-	-	-	-	-	-	++++



# Process problem

## ■ Negative eller manglende processor




Buitenwerf, R., Sandel, B., Normand, S. Mimet, A. & Svenning, J.-C. 2018. Land-surface greening suggests vigorous woody regrowth throughout European semi-natural vegetation. *Global Change Biology* 24:5789–5801.

Foto: JCS

## 2: Wildness as a value

"There are no words that can tell the hidden spirit of the **wilderness**, that can reveal its mystery, its melancholy, and its charm"

–Theodore Roosevelt,  
1858-1919






Yellowstone National Park


Foto: JCS



### 3: Shifting baseline



- Critically assess ecological integrity
  - Especially relative to evolutionary conditions
- Active restoration may be needed to not get locked into biodiversity-poor degraded persistent states
  - Especially as societally relevant time scales are decades, not millenia



### 3: Shifting baseline

# Rewilding as tool to increase our understanding of nature




Photo: Lascaux (www)


# 4: Scaling

- Functional area for nature/biodiversity is the biggest need to encounter the biodiversity crisis
- **Upscaling** crucial
  - Only practical & effective if strongly based on **autonomous** natural processes
    - Cost/labor
    - Value for biodiversity




Selfsown oak (*Quercus robur*)

## 5: Dynamic



**Fig. 7.** Time series of the percentage of the global land area ( $60^{\circ}\text{S}$ – $90^{\circ}\text{N}$ ) assigned different climate types compared to the present day condition (1961–90). The black (green) lines are the temporal variations based on the dataset from the University of Delaware (UD) and CPC, respectively. Thick blue (red) lines are the ensemble average of the 20 models from historical/RCP8.5 (RCP4.5) simulations. The gray shading denotes one standard deviation of the 20 models from historical/RCP8.5 simulations. Blue (light blue) dashed line is the temporal variation based merely on the temperature (precipitation) changes from historical/RCP8.5 simulations. Pink (pink dashed) line is the temporal variations based merely on the temperature (precipitation) changes from the UD dataset. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

# Comparison to past climate



Temperature anomalies are relative to 1961–1990 global means and are composited from five proxy-based reconstructions, modern observations, and future temperature projections for four emissions pathways. Pal, Paleocene; Mio, Miocene; Oli, Oligocene.

# Comparison to past climate



Ellesmere Island, 3.4 mio years ago

# **IMPLEMENTATION**

# Framework for design & implementation

- Design:
  - Trophic complexity
  - Natural disturbances
  - Dispersal/Connectivity
- Implementation
  - 1) Ecosystem status assessment
  - 2) Social-ecological constraints
  - 3) Adaptive management

RESEARCH

## REVIEW SUMMARY

### REWILDING

## Rewilding complex ecosystems

Andrea Perino\*, Henrique M. Pereira\*, Laetitia M. Navarro, Néstor Fernández, James M. Bullock, Silvia Ceaușu, Ainara Cortés-Aizanda, Roel van Klink, Tobias Kuehmerle, Angela Lomba, Guy Pe'er, Tobias Plieninger, José M. Rey Benayas, Christopher J. Sandom, Jens-Christian Svenning, Helen C. Wheeler

**BACKGROUND:** Rapid global change is creating fundamental challenges for the persistence of natural ecosystems and their biodiversity. Conservation efforts aimed at the protection of landscapes have had mixed success, and there is an increasing awareness that the long-term protection of biodiversity requires inclusion of flexible restoration along with protection. Rewilding is one such approach that has been both promoted and criticized in recent years. Proponents emphasize the potential of rewilding to tap opportunities for restoration while creating benefits for both ecosystems and societies. Critics discuss the lack of a consistent definition of rewilding and insufficient knowledge about its potential outcomes. Other criticisms arise from the mistaken notion that rewilding actions are planned without considering societal acceptability and benefits. Here, we present a framework for rewilding actions that can serve as a guideline for researchers and managers. The framework is applicable to a variety of rewilding approaches, ranging from passive to trophic rewilding, and aims

to promote beneficial interactions between society and nature.

**ADVANCES:** The concept of rewilding has evolved from its initial emphasis on protecting large, connected areas for large carnivore conservation to a process-oriented, dynamic approach. On the basis of concepts from resilience and complexity theory of social-ecological systems, we identify trophic complexity, stochastic disturbances, and dispersal as three critical components of natural ecosystem dynamics. We propose that the restoration of these processes, and their interactions, can lead to increased self-sustainability of ecosystems and should be at the core of rewilding actions. Building on these concepts, we develop a framework to design and evaluate rewilding plans. Alongside ecological restoration goals, our framework emphasizes people's perceptions and experiences of wildness and the regulating and material contributions from restoring nature. These societal aspects are important outcomes and may be critical factors for the success of


rewilding initiatives (see the figure). We further identify current societal constraints on rewilding and suggest actions to mitigate them.

**OUTLOOK:** The concept of rewilding challenges us to rethink the way we manage nature and to broaden our vision about how nature will respond to changes that society brings, both intentionally and unintentionally. The effects of rewilding actions will be specific to each ecosystem, and thus a deep understanding of the processes that shape ecosystems is critical to anticipate these effects and to take appropriate management actions. In addition, the decision of whether a rewilding approach is desirable should consider stakeholders' needs and expectations. To this end, structured restoration planning—based on participatory processes involving researchers, managers, and stakeholders—that includes monitoring and adaptive management can be used. With the recent designation of 2021–2030 as the “decade of ecosystem restoration” by the United Nations General Assembly, policy- and decision-makers could push rewilding topics to the forefront of discussions about how to reach post-2020 biodiversity goals. ■

The list of author affiliations is available in the full article online.  
\*Corresponding author. Email: andrea.perino@idiva.de (A.P.); hpereira@idiva.de (H.M.P.)  
Cite this article as: Perino et al. *Science* 364, eaav5570 (2019). DOI: 10.1126/science.aav5570



**TOMORROW'S EARTH**  
Read more articles online at [scim.ag/tomorrows.earth](http://scim.ag/tomorrows.earth)




**Rewilding actions and outcomes are framed by societal and ecological context.** Rewilding can be assessed by representing the state of ecosystems in a three-dimensional space where each dimension corresponds to an ecological process. The difference in volume between the restored (yellow pyramid) and the degraded ecosystem (orange pyramid) is a proxy for the effects of rewilding on the self-sustainability of

the ecosystem. The dashed line within the yellow pyramid represents the societal boundaries that determine to what extent ecological processes can be restored. Rewilding actions can help push societal boundaries toward the ecological potential (orange arrows) by promoting societal support and opportunities for people to experience the autonomy of ecological processes in enjoyable ways.


Perino et al., *Science* 364, 351 (2019) 26 April 2019

1 of 1


# Framework for measuring rewilding progress



# A trophic rewilding scale to guide terminology and management




# Socio-ecological context



# Why is rewilding needed?

- 1) Evolutionary perspective on biodiversity
- 2) Wildness as a value
- 3) Overcome shifting baselines
- 4) Scaling
- 5) Dynamic



# Thanks

svenning@bios.au.dk

 @JCSvenning

- Collaborators!
- Funding
  - VILLUM FONDEN  
(*VILLUM Investigator grant*)  
*"Biodiversity Dynamics in a Changing World"*
  - Carlsberg Foundation "Semper Ardens": *MegaFauna ecosystem ecology from the deep prehistory to a human-dominated future*  
(*MegaPast2Future*)
  - Innovation Fund Denmark



**BIOCHANGE**

CENTER FOR BIODIVERSITY DYNAMICS  
IN A CHANGING WORLD

VILLUM FONDEN



CARLSBERG FOUNDATION

**MegaPast2Future**

**MegaFauna ecosystem ecology from the  
deep prehistory to a human-dominated  
future**



Innovationsfonden